

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281097876>

Resistance against fruit tree canker in apple – Evaluation of Disease Symptoms, Histopathological and RNA-Seq Analyses in Different Cultivars, Genetic Variation of *Neonectria ditis*...

Book · August 2015

CITATIONS

0

READS

1,684

1 author:



[Marjan Ghasemkhani](#)

Swedish University of Agricultural Sciences

19 PUBLICATIONS 62 CITATIONS

SEE PROFILE

Resistance against fruit tree canker in apple

Evaluation of Disease Symptoms, Histopathological and
RNA-Seq Analyses in Different Cultivars, Genetic
Variation of *Neonectria ditissima*

Marjan Ghasemkhani

*Faculty of Landscape Architecture, Horticulture and Crop Production Science
Department of Plant Breeding
Alnarp*

Doctoral Thesis
Swedish University of Agricultural Sciences
Alnarp 2015

Acta Universitatis Agriculturae Sueciae

2015:77

Cover: Apple flowers (photo: T. Larsen), Fruit tree canker symptoms on an apple tree (photo: H. Nybom), Fruit tree canker symptoms on main trunk, Sporodochium of *Neonectria ditissima*, Fungal hyphae of *N. ditissima* in the cortex, Fungal hyphae of *N. ditissima* under periderm (last four photos: M. Ghasemkhani).

ISSN 1652-6880

ISBN (print version) 978-91-576-8352-6

ISBN (electronic version) 978-91-576-8353-3

© 2015 Marjan Ghasemkhani, Alnarp

Print: SLU Service/Repro, Alnarp 2015

Resistance against fruit tree canker in apple

Evaluation of Disease Symptoms, Histopathological and RNA-Seq Analyses
in Different Cultivars, Genetic Variation of *Neonectria ditissima*

Abstract

Neonectria ditissima (formerly *Neonectria galligena*, anamorph *Cylindrocarpon heteronema*) is the causal agent of fruit tree canker, which is regarded as a serious economic problem in apple production. This fungus is closely associated with mild and wet conditions, and climate therefore has an important impact on its geographic distribution. Notable damage to apple trees is especially common in, e.g., North-Western Europe. Complete resistance has not yet been reported. In addition, no major genes have as yet been identified and resistance appears to be controlled mainly by additive gene action.

Apple cultivars varied significantly in level of partial resistance towards apple canker after artificial inoculation of detached shoots and/or potted trees with macroconidia of *N. ditissima*. The inter-specific crabapple hybrid 'Prairifire' showed the least damage, indicating that valuable sources of resistance can be found in *Malus* species. *Neonectria ditissima* could be detected in artificially infected apple trees using a species-specific β -tubulin gene-based qPCR assay two months after inoculation. When quantified in different cultivars, a positive relationship between *N. ditissima* biomass and the size of canker-induced lesions was found. Interaction between plant and pathogen was studied using light microscopy of apple woody tissue; fungal hyphae were found in all cell types, and gel formation was observed in both a susceptible and a resistant cultivar but infection appears to progress more rapidly in the susceptible cultivar. Genetic variation among different *N. ditissima* isolates was investigated using SSR and AFLP markers. Comparatively low levels of genetic differentiation among orchards indicate that gene flow is prominent. Analyses of single-ascospore isolates from the same perithecia suggest that this fungus is heterothallic and therefore outcrossing. Furthermore, a focused genomics approach was applied in order to identify differentially expressed genes in response to fungal attack on apple trees. Obtained data indicate that apple cultivars inoculated with *N. ditissima* exhibit significant upregulation of defence-related genes and genes involved in detoxification, peroxidase-related reactions, phenylpropanoid metabolism and the lignification process.

Keywords: anatomy, β -tubulin gene, defence mechanism, genetic variation, mating system, qPCR, RNA-Seq, transcriptome

Author's address: Marjan Ghasemkhani, SLU, Department of Plant Breeding,
P.O. Box 101, 230 53 Alnarp, Sweden
E-mail: Marjan.Ghasemkhani@slu.se

Dedication

To my parents for their endless love, support and encouragement.

To my husband for his patience, love, friendship and humour.

Absence of understanding does not warrant absence of existence.

Ibn Sina

Contents

List of Publications	7
Abbreviations	11
1 Background	13
1.1 Apple (<i>Malus x domestica</i>)	13
1.1.1 History and origin	13
1.2 Fruit tree canker (<i>Neonectria ditissima</i>)	14
1.2.1 History and origin	14
1.2.2 Geographic distribution	15
1.2.3 Morphology	15
1.2.4 Growth of <i>N. ditissima</i> in apple tree tissues	18
1.2.5 Life cycle and infection	18
1.2.6 Symptoms	20
1.3 Management of the fruit tree canker	20
1.3.1 Development of healthy apple cultivars	21
2 Aim and objectives	23
3 Material and methods	25
3.1 Plant and fungal materials	25
3.2 Estimation of resistance towards <i>N. ditissima</i> in apple cultivars	26
3.2.1 Inoculation procedure	26
3.3 Detection and quantification of <i>N. ditissima</i> with quantitative real-time PCR	28
3.3.1 Primer design and validation of primer specificity	28
3.3.2 Experimental procedures	29
3.4 Histopathology of canker on <i>Malus</i> caused by <i>N. ditissima</i>	29
3.4.1 Experimental procedures	29
3.4.2 Data evaluation	30
3.5 Genetic variation in <i>N. ditissima</i>	30
3.5.1 DNA extraction and amplification	30
3.5.2 Data analysis	30
3.6 Gene expression analysis using RNA-sequencing data	32
3.6.1 Experimental procedures	32
3.6.2 Data analysis	32

4	Results and discussion	35
4.1	Variation among apple cultivars inoculated with spores of <i>N. ditissima</i>	35
4.2	Quantitative real-time PCR for detection and quantification of <i>N. ditissima</i> in susceptible and resistant apple cultivars	38
4.3	Histopathology of infection and colonization of susceptible and resistant apple cultivars by <i>N. ditissima</i>	39
4.4	Genetic diversity and genetic relationships of <i>N. ditissima</i>	42
4.5	Identification of differentially expressed genes in partially resistant and susceptible cultivars in response to inoculation with <i>N. ditissima</i>	44
5	Conclusions and future perspectives	49
	References	51
	Acknowledgements	63

List of Publications

This thesis is based on the work contained in the following papers, referred to by Roman numerals in the text:

- I **Ghasemkhani M.**, Liljeroth E., Sehic J., Zborowska A. and Nybom, H. (2015). Cut-off shoots method for estimation of partial resistance in apple cultivars to fruit tree canker caused by *Neonectria ditissima*. *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science*, 65(5): 412-421.
- II Garkava-Gustavsson L., **Ghasemkhani M.**, Zborowska A., Englund J.-E., Lateur M. and van de Weg E. Approaches for evaluation of resistance to European canker (*Neonectria ditissima*) in apple. *Acta Hortic.* (In press).
- III **Ghasemkhani M.**, Holefors A., Marttila S., Zborowska A., Rur M., Rees-George J., Nybom H., Everett K., Scheper R. and Garkava-Gustavsson L. Detection and quantification of *Neonectria ditissima* in apple trees by real-time quantitative PCR and visualization by microscopy. (Submitted).
- IV **Ghasemkhani M.**, Garkava-Gustavsson L., Liljeroth E. and Nybom H. Assessment of diversity and genetic relationships of *Neonectria ditissima*: the causal agent of fruit tree canker. (Manuscript).
- V **Ghasemkhani M.**, Canbäck B., Alexandersson E., Nybom H. and Garkava-Gustavsson L. Identification of differentially expressed genes associated with response to fruit tree canker in apple using RNA-Seq analysis. (Manuscript).

Papers I and II are reproduced with the permission of the publishers.

The contribution of Marjan Ghasemkhani to the papers included in this thesis was as follows:

- I Planned and performed some of the experimental work together with co-authors, performed all statistical analyses and wrote the manuscript.
- II Performed the experimental work together with co-authors and participated in writing of the manuscript.
- III Planned the experiments together with co-authors. Performed a large part of the experimental work, performed all statistical analyses and wrote the manuscript.
- IV Planned the experiments together with co-authors. Performed all the experimental work, evaluated and analysed all data and wrote the manuscript.
- V Planned the experiments together with co-authors. Performed all the experimental work, evaluated and analysed the data together with one of the co-authors (bioinformatician) and wrote the manuscript.

Related articles by Marjan Ghasemkhani but not included in this thesis:

- I. **Ghasemkhani M.**, Sehic J., Ahmadi-Afzadi M., Nybom H. and L. Garkava-Gustavsson. Screening for partial resistance to fruit tree canker in apple cultivars. *Acta Hortic.* (In press).
- II. **Ghasemkhani M.**, Holefors A., Zborowska A., Scheper R., Everett K., Nybom H and Garkava-Gustavsson L. Development of a qPCR detection procedure of fruit tree canker caused by *Neonectria ditissima*. *Acta Hortic.* (In press).
- III. Nybom H., Røen Dag., Karhu S., Garkava-Gustavsson L., Tahir I., Haikonen T., Røen K., Ahmadi-Afzadi M., **Ghasemkhani M.** and S. H. Hjeltnes. Prebreeding for future challenges in Nordic apples; susceptibility to fruit tree canker and storage diseases. *Acta Hortic.* (In press).

Abbreviations

AMOVA	Analysis of molecular variance
ANOVA	Analysis of variance
AUDPC	Area under the disease-progression curve
BC	Christian calendar concepts
CYPs	Cytochromes P450
dai	days after inoculation
DEGs	Differentially expressed genes
<i>ELI3-1</i>	Elicitor-activated gene 3-1
gDNA	Genomic DNA
ITS	Internal transcribed spacer
KEGG	Kyoto encyclopedia of genes and genomes
LOD	Limit of detection
LSD	Fisher's least significant difference
MEA	Malt extract agar
<i>MLO12</i>	Mildew resistance locus O 12
mRNA	Messenger RNA
PCA	Principal component analysis
PCR	Polymerase chain reaction
PR	Pathogenesis-related
qPCR	Quantitative real-time polymerase chain reaction
QTL	Quantitative trait loci
RNA-Seq	RNA sequencing

1 Background

1.1 Apple (*Malus × domestica*)

Apple, *Malus × domestica* (L.) Borkh., is a member of the Rosaceae family. It is the fourth most widely grown fruit in the world after citrus, grapes and banana (O'Rourke, 2003; Khachatourians, 2002), and is cultivated in temperate and subtropical areas. Due to its wide accessibility, apple is also one of the most popular and widely consumed deciduous fruits. Nearly half of the production is consumed as fresh fruit and most of the remainder is processed into apple juice, apple jam, apple butter and canned apple sauce. Charoset (apple relish) and dehydrated apple slices are other important commercial products. From a medicinal point of view, fresh apple has been reported to act as a purgative, and to reduce incidence of dental caries, prevent constipation, supply extra energy for heavy exercise and help to control obesity (Bakhru, 1995).

1.1.1 History and origin

Historical records and prehistoric remains indicate the existence of cultivation and dispersal of apple in Asia and Europe more than several thousand years ago. Apples were widely cultivated in the near East almost 3000 BC (e.g., Turkey, Syria, Iraq) (Hancock, 2008) and in the Persian Empire (e.g., Iran and beyond) by 500 BC (Hancock, 2008; Luby *et al.*, 2003). The cultivation of apple subsequently spread through Greece and the Roman Empire to Europe. Apple trees were soon planted in almost the whole of Europe and a large number of different cultivars were identified by the end of the 18th century (Hancock, 2008; Luby *et al.*, 2003; Juniper *et al.*, 1996).

Using molecular markers, Harris *et al.* (2002) showed that the wild apple *Malus sieversii*, located in Central Asia, is the major contributor to the domesticated apple. Other *Malus* species have also contributed to the gene pool, both in ancient times and more recently as donors of resistance to pests

and diseases. Almost all commonly grown apple cultivars are treated as *M. × domestica* but some crabapples (*Malus* spp.) are still grown, mainly for ornamental purposes and as pollen donors in commercial orchards.

Today, the commercial apple production exceeds 70 million metric tons in the world, with China, U.S., Turkey, Poland, India, Italy, Iran, Chile, France and Russia being the leading producers (FAO, 2013; O'Rourke, 2003).

1.2 Fruit tree canker (*Neonectria ditissima*)

Several different fungi produce cankers on apple trees and reduce both yield and growth, and may ultimately lead to death of the trees. Fruit tree canker, caused by the fungus *Neonectria ditissima* (Tul. & C. Tul.) Samuels & Rossman (*Neonectria galligena* (Bres.) Rossman & Samuels formerly known as *Nectria galligena*), is one of the most important diseases of apple trees and is regarded as a serious economic problem, especially in North-Western Europe (Weber, 2014; Rossman & Palm-Hernandez, 2008; Rossman *et al.*, 1999).

1.2.1 History and origin

The first reports on *Neonectria ditissima* probably date from approximately 1880 when two studies of putative fruit tree canker were published; Goethe (1880) and Hartig (1880) reported of a similar disease on a selection of broad-leaved trees, especially copper beach. The early stages of canker formation on apple trees were described by Wiltshire (1921), who reported that *N. ditissima* can enter through cracks, which occur in the spring when the buds are swelling, after leaf-fall in autumn and through lesions produced by the scab fungus *Venturia inaequalis*. A detailed description of the disease and the anatomy of the lesions was provided by Zeller (1926), while Cayley (1921) described the life-history of *N. ditissima* in order to facilitate the use of fungicides during the spore-producing periods.

Apple canker, *Neonectria* canker, and European canker are common names of fruit tree canker. A previous study using RAPD and ribosomal DNA polymorphisms has shown that this pathogen originated in North America (Plante *et al.*, 2002). *Neonectria ditissima* belongs to the phylum *Ascomycota*, class *Sordariomycetes*, subclass *Hypocreomycetidae* and order *Hypocreales*. According to evidence from molecular phylogenetic studies and information about the asexual and sexual reproduction of *N. ditissima*, this pathogen belongs to family *Nectriaceae*, and *Neonectria* is the only identified and proven teleomorph of *Cylindrocarpon heteronema* (Chaverri *et al.*, 2011; Castlebury *et al.*, 2006). During the past 150 years, the scientific name of this

disease has changed frequently and currently, *N. ditissima* (Tul. & C. Tul.) Samuels & Rossman has taken precedence over *N. galligena* (Rossman & Palm-Hernandez, 2008; Castlebury *et al.*, 2006).

1.2.2 Geographic distribution

Fruit tree canker has been reported from Australia, Chile, Europe, South Africa, New Zealand and Japan (Weber, 2014; Beresford & Kim, 2011; Xu & Robinson, 2010; Grove, 1990). A minimum number of days with rainfall (> 30 %) and five months per year with specific temperatures (> 8 h at 11–16 °C) are required for this pathogen to cause severe canker (Beresford & Kim, 2011). Local climatic factors have a perceptible impact on the distribution and severity of canker. Severe canker frequently occurs on trees growing on poorly drained soils, on soils that have pockets that are poorly drained at lower altitudes and on infertile soils at higher altitudes (Weber, 2014; Brandt, 1964). *Neonectria ditissima* infects apple, pear (usually less severely than apple), and many hardwood forest trees such as beech, birch, maple, aspen, quince and hickory (Braun & Craig, 2008; Farr *et al.*, 1989; Flack & Swinburne, 1977).

1.2.3 Morphology

Morphological characteristics of *N. ditissima* are observed on asexual reproduction organs, i.e., sporodochia, conidiophores and conidia, and sexual reproduction organs, i.e., perithecia, asci and ascospores. Sporodochia consist of conidiophores (Figure 1), which are usually branched and arise from the white mycelium on the surface of the bark (Figure 2) (Agrios, 1997).




Figure 1. Sporodochia of *Neonectria ditissima*.

The asexual stage of this pathogen, *Cylindrocarpon heteronema*, produces two kinds of conidia; macroconidia and microconidia. Fully developed mycelia produce sporodochia on the surface of infected and dead bark (Agrios, 1997). Macroconidia are ellipsoidal or cylindrical with rounded ends, and are produced at the end of the conidiophores of these sporodochia (Figure 3). Microconidia are instead single-celled and vary in shape and size (Ghasemkhani, 2012). Macroconidia are extremely infectious, while the epidemiological role of microconidia is unclear (Weber, 2014; Saure, 1961; Zeller, 1926).

Perithecia, the sexual fruiting body of this fungus, are produced on infected tissue. They are bright red and vary in size, depending on environmental conditions and position on the bark. Perithecia can be found much later in the year than sporodochia (Figure 4) (Weber, 2014; Lortie, 1964).




Figure 2. Conidiophores from a sporodochium produced on apple bark, A: part of conidiophore, B: densely aggregated mass of conidiophores.

The asci of *N. ditissima* are cylindrical and contain eight ascospores (Figure 5A) (Ainsworth & Bisby, 2011; Hanlin, 1971). Ascospores are two-celled, one-septate and ellipsoidal, and vary moderately in shape and size (Figure 5B). They are released in spring and early summer but there is also a short period of discharge in the autumn, and can spread over long distances (Swinburne, 1971). Weather conditions, especially humidity, have a significant impact and it has been reported that ascospore discharge does not occur unless the leaves are wet (Wessel, 1979; Lortie & Kuntz, 1963; Wiltshire, 1921).




Figure 3. Macroconidia of *Neonectria ditissima* collected on the bark of an apple tree.



Figure 4. Perithecia of *Neonectria ditissima*, produced during winter. Photo: Bruce A. Watt.




Figure 5. *Neonectria ditissima* produces asci in perithecia, A: two-celled ascospores, B: one ascus with 8 two-celled ascospores. Photo: Bruce A. Watt.

1.2.4 Growth of *N. ditissima* in apple tree tissues

According to Crowdy (1949), hyphae of *N. ditissima* attack xylem vessels, tracheids and fibers, and have also been detected inside xylem vessels (Langrell, 2000). The fungal hyphae are abundant close to the infection sites and become less frequent further away from these sites (Sakamoto *et al.*, 2004). The fungus also penetrates to the soft tissues outside the xylem. Peripheral tissue can be stimulated to form a phellogen barrier, which blocks the vascular tissue and helps to restrict further infection (Clatterbuck, 2006). This barrier can break down if attacked by a large aggregate of mycelia, possibly due to mechanical pressure or to a high concentration of toxins produced by the fungus. However, the host soon forms a new phellogen barrier and this process is continuously repeated (Zeller, 1926). The fungus can also penetrate to the phloem fibers after first killing the living cells by secretion of toxin (Crowdy, 1949). The mycelia grow within cells at the early stage of infection and move between cells through pits. Later, the mycelia develop intercellularly and hyphae grow spirally around small groups of fibers within the bundles (Crowdy, 1949; Zeller, 1926).

1.2.5 Life cycle and infection

Neonectria ditissima survives in cankered tissues as mycelium and perithecia during the winter and/or during other detrimental conditions. Primary infection usually originates from ascospores in the spring (McCracken *et al.*, 2003; Welch, 1934). When the temperature is well above freezing and there is sufficient moisture, sporodochia are generally produced in the first year of new canker wounds. Perithecia usually develop and appear in late summer and autumn (Figure 6). Therefore, inoculum that can cause infection is available the whole year around except for during freezing conditions in winter or extremely dry periods in summer (Weber, 2014; Xu & Butt, 1994). The spores produced by the fungus are dispersed within the same tree and to neighbouring trees, with peak production of both kinds of spores at 10–16 °C in the autumn (Beresford & Kim, 2011; Latorre *et al.*, 2002).

There is, however, considerable climate-dependent variation in the reproduction and spread of this pathogen. In Chile and United States perithecia are rarely formed, and the principal source of inoculum is probably conidia (Dubin & English, 1970; Crowdy, 1952). In northern Germany, severe infection may result when the perithecia are sufficiently mature to release ascospores during leaf fall in the autumn (Weber, 2014). In the British Isles, most of the ascospores are instead discharged during spring (Weber, 2014; Swinburne, 1971; Munson, 1939).




Figure 6. The life cycle of fruit tree canker caused by *N. ditissima*. Figure modified from Agrios (1997).

Dissemination of conidia and ascospores depends on e.g., climate conditions such as rainfall during the season and rain splash, wind, and the activity of insects, all of which are considered as carriers. Leaf scars in the autumn constitute the most important entry sites for infection (Swinburne *et al.*, 1975; Saure, 1961; Wiltshire, 1921). Other wounds e.g., cracks, frost injuries, pruning wounds in winter and summer, wounds caused by insects or by the picking of fruits, and apple scab lesions can also act as entry sites for infection (Weber, 2014; Pijut, 2006; Mols & Boers, 2001; Dewey & Swinburne, 1995; Swinburne, 1971; Dubin & English, 1970; Kennel, 1963; Saure, 1961). Different types of wounds, i.e. natural or artificial, are available on trees throughout the whole year, and the spores can grow and germinate instantly, and establish in the bark tissue within 3 to 4 hours (Xu *et al.*, 1998; Brandt, 1964). Conidia can spread during the growing season to other trees and this cycle is repeated several times.

A previous study showed that 50 and 5000 macroconidia per leaf scar, caused 20% and 90% infection, respectively (Dubin & English, 1970) while 100 macroconidia on fresh pruning wounds are needed to cause 32% infection (Xu *et al.*, 1998). Sometimes, wounds are not sufficiently deep for successful

penetration by the fungus, and the infection is limited to the cortex (Wiltshire, 1921). After initial entry of *N. ditissima* and establishment of infection, wound callus grows around the infected area in order to prevent the spread of the infection into new tissue.

1.2.6 Symptoms

Neonectria ditissima can infect trees of all ages. The first symptoms of infection on apple branches and stems consist of small dark reddish spots in an area surrounding the infection site. Sunken areas can develop around the infected sites, e.g., on shoot bases, leaf scars and wounds (Figure 7A) (Crowdy, 1949).

Large lesions on main branches or stems usually originate from infection at the base of side shoots that have extended into the larger branches. The host forms phellogen barriers around the infected sites in response to infection resulting in swelling area (Beltra *et al.*, 1969). Concentric ridges are observed in exposed wood of old cankers, caused by differences in the seasonal growth rate of the fungus and the host (Figure 7B) (Crowdy, 1949). Cross-sections of infected branches mostly show a browning area within the wood, and hyphae of *N. ditissima* can be detected in the lumen of xylem vessels (Weber, 2014; Ventura *et al.*, 2004; Dewey & Swinburne, 1995). Conidial spore masses, looking like white fruiting bodies, can be identified on young cankered shoots in the summer and early autumn, while perithecia are recognized as red fruiting bodies in spring, autumn and winter (Weber, 2014; Crowdy, 1949).

Neonectria ditissima can also cause fruit rotting, but symptoms are seldom observed before harvest. During storage, symptoms can develop on all parts of the fruit, and the skin over the rotted area becomes dark brown (Weber, 2014).

1.3 Management of the fruit tree canker

Control of fruit tree canker is difficult since the fungus produces spores the whole year and there are always suitable entry sites on apple trees for infection. A combination of methods, including orchard management procedures, chemical treatments and choice of comparatively resistant cultivars, is required to keep the disease under at least partial control. These procedures include cutting out all infected tissue and covering the wounds with disinfectant as soon as an infected tree has been detected during winter pruning. A number of chemicals like Bordeaux paste, Copper-Count-N, and Cuprofix Ultra 40 have been used on the cankers for damage control (Cooke, 1999; English *et al.*, 1979; Swinburne *et al.*, 1975), as also spraying of the trees with e.g., copper oxychloride (Cooke *et al.*, 1993) but these chemicals are not permitted any

longer in an increasing number of countries. Scaniavital Kambium, which is a Swedish product made from natural sea bottom-derived paste containing silica, calcium and various organic components, has been commercialized for treatment of fruit tree canker (www.nordiskalkali.se). Unfortunately, none of these protectant procedures are sufficient, and it becomes especially difficult to control the disease if it is established already in young trees in newly planted orchards (Cooke, 1999; Swinburne *et al.*, 1975).



Figure 7. A: Sunken area around initial symptoms on young branches infected by *N. ditissima*, B: concentric rings typical of an older canker infection.

1.3.1 Development of healthy apple cultivars

It is well known that both *Malus* species and apple cultivars show variable levels of resistance to fruit tree canker (Sasnauskas *et al.*, 2006; Kozlovskaya *et al.*, 1999; Lateur & Populer, 1994; van de Weg, 1989), and some of the traditional cider apple cultivars are comparatively resistant as also some rootstocks like ‘M.1’ and ‘M.12’ (Moore, 1960). However, complete resistance has not yet been reported (Ghasemkhani *et al.*, 2015; Garkava-Gustavsson *et al.*, 2013; van de Weg, 1989; van de Weg, 1987) and there are few plant breeding programs focused on developing apple genotypes with high levels of resistance to this pathogen.

Level of resistance is generally associated with the mode of inheritance. Single-gene or oligogenic resistance is controlled by one or a few major-effect genes (also known as qualitative or vertical resistance) while polygenic resistance is controlled by many genes each of which have a restricted phenotypic effect (quantitative or horizontal resistance) (Balconi *et al.*, 2012;

Baker & Cook, 1974). In qualitative resistance, most of the determined R gene(s) are dominantly inherited but this can vary with pathogen isolate (Roelfs, 1988), disease phenotype (Johnson, 1992), genetic background or environment (Balconi *et al.*, 2012). These genes are often race-specific, providing full protection until there is a breakdown in resistance, as has happened with the *Vf*-gene against apple scab (Bénaouf & Parisi, 2000; Parisi *et al.*, 1993). However, specific R-gene resistance has not been observed against fruit tree canker.

By contrast, quantitative resistance usually provides only partial resistance (Yang *et al.*, 1997). The roles of specific loci in such traits can be investigated by identification and genetic mapping of QTL (quantitative trait loci) (Balconi *et al.*, 2012; Talukder *et al.*, 2004). Possibly a limited number of major QTLs is involved in the resistance to apple canker (van de Weg & Jansen, 1989), and a more recent study has demonstrated that resistance to apple canker is controlled predominantly by additive gene action (Gelvonauskiene *et al.*, 2007).

2 Aim and objectives

Most of the studies have been conducted on the epidemiology of this disease and there are few studies on host plant resistance while other aspects of the resistance remain unclear. In addition, the mechanism of resistance in apple cultivars against fruit tree canker is poorly understood. In this thesis work, susceptibility of different apple cultivars to fruit tree canker was investigated. In addition, studies on detection, quantification of this fungus and plant-pathogen interaction in artificially infected trees and genetic variation of the pathogen, were performed. The last step was to elucidate how the apple defence is affected by this pathogen on a molecular level. The more specific objectives were to:

- Determine levels of resistance to fruit tree canker in an apple germplasm collection using *N. ditissima* spore inoculation with two different plant models: detached shoots and potted trees, respectively.
- Detect and quantify biomass of *N. ditissima* in cankered apple cultivars and analyse the association between pathogen DNA accumulation in different apple cultivars and levels of disease severity.
- Investigate the ultrastructure of plant-pathogen interaction by microscopy to improve understanding of partial resistance to *N. ditissima*.
- Evaluate genetic variation of *N. ditissima* among and within apple orchards and investigate the mating system using SSR and AFLP markers.

- Apply RNA-Seq analysis to identify differentially expressed genes that can help to elucidate the mechanisms behind resistance towards *N. ditissima* in apple.

3 Material and methods

3.1 Plant and fungal materials

Apple cultivars, which are commonly grown in the Nordic countries and have a potential interest in breeding programs, have been used together with a few international and/or in other ways special cultivars.

For the detached-shoots spore-inoculation experiments, dormant shoots were collected from trees in the apple germplasm orchard at Balsgård (N 56°6', E 14°9'), Swedish University of Agricultural Sciences, in southern Sweden, and from one cultivar in Finland. The shoots were kept in cold storage (+4 °C) from collection to four days prior to inoculation and no fungicide was applied.

For the potted-trees spore-inoculation experiments, one-year-old apple trees grafted on the rootstock M9, were produced in a Swedish nursery using budwood from Balsgård. The potted trees were kept in an unheated greenhouse and watered weekly. No fungicides and fertilizers were used.

For qPCR experiments, histopathology and gene expression study, one-year-old trees were produced and handled as previously described.

For the genetic variation study, single-ascospore isolates were produced from perithecia taken from naturally occurring cankers on apple branches collected in seven orchards in southern Sweden and in one orchard in Belgium (Table 1).

In order to prepare the spore suspension for spore-inoculation experiments, sporodochia of *N. ditissima* from naturally occurring cankers on apple branches were collected in the orchards at Balsgård. Macroconidia of *N. ditissima* were distinguished based on morphological characters, and a macroconidial suspension was obtained. The concentration was adjusted to 1×10^5 conidia mL⁻¹.

For the qPCR and gene expression studies, the macroconidial suspension, as previously described, was grown on agar media (1.5%). Next day, single

macroconidia were cultured on agar media (1.5%) to produce sporodochia. Later, produced sporodochia were used for spore suspensions.

Table 1. Geographic origin of isolates of *N. ditissima* used in this study

Number of isolates	Orchard	Latitude and longitude	Year
3	Jonstorp, Sweden	(N 56°13', E 12°40')	2013
2	Julita, Sweden	(N 59°11', E 16°1')	2013
2	Jönköping, Sweden	(N 57°46' N, E 14°9')	2013
2	Kivik, Sweden	(N 55°41', E 14°13')	2013
26	Balsgård, Sweden	(N 56°6', E 14°9')	2013
3	Bjärred, Sweden	(N 55°43', E 13°1')	2013
4	Gembloux, Belgium	(N 54°34', E 4°42')	2014
2	Stockholm, Sweden	(N 59°19', E 18°4')	2014

Isolator: Marjan Ghasemkhani

3.2 Estimation of resistance towards *N. ditissima* in apple cultivars

In the first study (Paper I), genetic differences in resistance to fruit tree canker was investigated in 30 apple cultivars, which were screened using wound-inoculated cut shoots in a standard greenhouse across two years (2012 and 2013). Eleven of these apple cultivars were also inoculated in a greenhouse with advanced climate control (biotron) in order to evaluate the robustness of the method. In the second study (Paper II), the correspondence of the results on resistance/susceptibility between years and between two plant models (cut shoots and one year-old trees) was verified on 15 cultivars.

3.2.1 Inoculation procedure

In the first experiment, shoots were transferred to a standard greenhouse at Balsgård and a greenhouse with advanced climate control at Alnarp (biotron) four days before inoculation, and were placed in 1L glass bottles. The glass bottles were kept under a plastic tent at Balsgård (Figure 8A) and in a chamber at Alnarp. Three axillary buds on each shoot were inoculated with spore suspension. A white petroleum jelly was used to cover the wounds for protection immediately after inoculation (Figure 8B), and was removed with tissue paper four days later.

In a second study, one-year-old trees and cut shoots were kept in an unheated greenhouse and in the biotron at Alnarp, respectively. They were inoculated as previously described. For evaluation of infection percentage,

another set of one-year-old trees were kept in the field during natural leaf fall. These trees were subjected to overhead irrigation at a precipitation regime normal for a Swedish autumn (20 mm/day). More details on the procedures are provided in Papers I and II.

3.2.2 Data evaluation and statistical analysis

In the first study (Paper I), lesion length was measured with a digital calliper at five time points at specified time intervals: five days in both years in the standard greenhouse and in the biotron 2013; and seven days in the biotron in 2012. In total, 90 and 120 evaluations were obtained in the standard greenhouse and in the biotron (3 and 4 shoots \times 2 replicates in time \times 3 leaf scars/shoot \times 5 assessments), respectively, for each cultivar in each year.




Figure 8. A: Cut-off apple shoots under a plastic tent in a standard greenhouse at Balsgård, B: The wounds were covered with white petroleum jelly just after inoculation.

Level of partial resistance was quantified in three different ways for the 11 cultivars assessed in both facilities: infection percentage, incubation period and area under the disease-progression curve (AUDPC). Separate two-way ANOVAs were calculated for these three variables with effects attributed to cultivar, year, replicate and facility. In order to investigate correspondence between replicates in both facilities, Pearson correlation tests were performed. A set of one-way ANOVAs was calculated for the 30 cultivars in both years in the standard greenhouse using AUDPC values, and an LSD post-hoc test was

used to compare the cultivars. In addition, Pearson correlation tests were applied to examine correspondence of the results between years.

In the second study (Paper II) one-way ANOVA was calculated for each plant model (cut shoots and potted trees) separately. A two-way ANOVA was performed with AUDPC as a dependent factor, and cultivar and year as independent variables for both plant models, and a Tukey's test was used to compare the cultivars. Pearson correlation tests were applied to investigate relationships between two plant models across two years with AUDPC values. Infection percentage of one-year-old trees was evaluated by dividing the proportion of infected scars to the total number of leaf scars. Spearman's rank correlation test was carried out to investigate possible relationships between the two years for each model, and between AUDPC and infection percentage. Calculations were conducted using SAS version 9.3, Minitab version 16 and Excel.

3.3 Detection and quantification of *N. ditissima* with quantitative real-time PCR

In this part of the thesis (Paper III), specificity of newly designed primer pairs was investigated. One primer pair, which passed several validation tests, was subsequently used to quantify the biomass of *N. ditissima* in cankered trees of artificially infected apple cultivars.

3.3.1 Primer design and validation of primer specificity

The primer pairs were designed from a β -tubulin gene by aligning regions conserved within *N. ditissima* with sequence information of different fungi that were deposited in GeneBank. Several primer combinations were compared using *in silico* analysis. Two primer sets were used for future evaluation.

Genomic DNA of *N. ditissima* was amplified with the designed primer pairs and resulting PCR products were purified, sequenced and assembled. Assembled sequences were aligned to gDNA sequences of *N. ditissima*. In the next step, extracted DNAs of *N. ditissima* from different isolates (Table 2 in Paper III) and different fungal pathogens that have been found on apple wood (Gomori, 1955) (Table 1 in Paper III), were used to evaluate primer specificity evidenced by presence or absence of a single PCR product of gDNA of *N. ditissima*. And finally, the melting curves with a single high peak during qPCR assay were evaluated. This method was also used for detection of *N. ditissima* and in investigations of the association between pathogen DNA accumulation and lesion size of infected apple cultivars.

The slope of a standard curve was used to estimate the amplification efficiency of the best primer pair using 5-fold serial dilution of gDNA of *N. ditissima*, and limit of detection (LOD) of *N. ditissima* was determined by running a total of 60 samples.

3.3.2 Experimental procedures

For quantification of *N. ditissima* in infected apple cultivars, one-year-old apple trees representing six cultivars, both susceptible and partially resistant, with three replications (trees) of each cultivar, in total 18 trees, were kept in an unheated greenhouse and inoculated with a macroconidial suspension. The samples were taken at two different time points; one month after inoculation and two months after inoculation, both at the point of infection and in the area surrounding the point of infection. DNAs were extracted and a qPCR assay was carried out. The reactions were run in 20 µL using 0.2 ml 96-well PCR plates. For amplification of *N. ditissima* and apple DNA, designed primer pairs and ubiquitin gene primers described by Botton *et al.* (2008) were used, respectively. More details on the procedures are provided in Paper III.

3.3.3 Data evaluation and statistical analysis

Analysis of qPCR data was carried out by Bio-Rad CFX manager Software v. 3.0.1. Pearson correlation was used to assess the relationship between lesion size and relative *N. ditissima* DNA quantity across the six cultivars. Statistical analyses were performed using the software R (R Development Core Team, 2013).

3.4 Histopathology of canker on *Malus* caused by *N. ditissima*

In order to understand how apple cultivars may defend themselves against fruit tree canker, plant-pathogen interaction in one susceptible and one partially resistant cultivar was compared.

3.4.1 Experimental procedures

A light microscopy study was undertaken on ‘Cox’s Orange Pippin’ chosen as a susceptible cultivar, and on ‘Santana’ chosen as a partially resistant cultivar. One-year-old potted trees and cut shoots were inoculated and sampled at the infection site, at 110 days after inoculation (dai) (trees), and at 5, 18, 29, and 36 dai (cut shoots).

Specimens were soaked in fixative (2% (w/v) paraformaldehyde, 2.5% (v/v) glutaraldehyde in 0.1 M Na-phosphate buffer, pH 7.2) during gentle agitation. The fixed samples were dehydrated and embedded in Spurr’s resin. Semithin

sections were cut and placed on object slides for light microscopy. Toluidine blue staining was used to observe the general structure and fungal hyphae. More details on the technical procedures are provided in Paper III.

3.4.2 Data evaluation

Sections were examined by a Leica DMLB light microscope and after determining the location of interest, photographs were taken with a Leica DFC450 digital camera.

3.5 Genetic variation in *N. ditissima*

In this part of the thesis, genetic diversity of *N. ditissima* was evaluated using single-ascospore isolates from different orchards (Paper IV). Single-ascospore isolates derived from the same perithecium were used for conclusions about mating system of this pathogen.

3.5.1 DNA extraction and amplification

Single-ascospore isolates were obtained from perithecia on pieces of bark and wood collected from different orchards in Sweden and Belgium (Table 2). In order to extract DNA of these isolates, fungal cultures on MEA 2% (Figure 9) were transferred into malt extract/glucose/peptone (MGP) liquid medium (Shaner & Finney, 1977) and incubated for 10 days (Figure 10). The mycelium was harvested, washed with dH₂O and freeze-dried. DNAs were extracted and quantity and quality checked.

Seven SSR primer pairs (Marra & Corwin, 2009), were used for SSR analysis. Genomic DNA was amplified using PCR.

AFLP analysis of DNA was carried out with an AFLP Microbial Fingerprinting kit. Eleven primer combinations were used. Reproducibility of AFLP amplification was checked by adding 5–8 randomly taken samples to each 96-well plate.

PCR products were separated on an ABI 3130xl DNA analyser and software package Gene-Marker v 1.85 was used for the fragment analysis. More details about experimental procedures are provided in Paper IV.

3.5.2 Data analysis

Genetic diversity parameters i.e., number of polymorphic loci (NPL), Nei's gene diversity (H), percentage of polymorphic loci (PPL) and Shannon's information index (I) were evaluated. Analysis of molecular variance (AMOVA) was performed in order to partition genetic variation into within- and among population components. Pairwise genetic similarities between

single-ascospore isolates obtained from the same or from different perithecia were estimated using Jaccard's similarity coefficients (Weising *et al.*, 2005). Similarity among isolates from the same or from different orchards was illustrated with a dendrogram constructed with UPGMA and a principal component analysis (PCoA). Mantel tests (Mantel, 1967) were performed in order to check the goodness-of-fit of the cluster analysis with the matrix on which it was based, and to investigate possible correlation between two genetic distance matrices based on SSR and AFLP markers, respectively. Calculations were conducted using POPGENE v 1.32 (Yeh *et al.*, 1997), NTSYS 2.02 (Tamura *et al.*, 2007) and Arlequin 3.11 (Excoffier *et al.*, 2005). More details about statistical calculations are provided in Paper IV.




Figure 9. *Neonectria ditissima*, culture on malt extract agar 2%. A, B, C: Fungal cultures with a circular mycelium. D: Erode margin of a fungal culture.



Figure 10. Different isolates of *Neonectria ditissima* growing in liquid media.

3.6 Gene expression analysis using RNA-sequencing data

Differentially expressed genes, that could be relevant for resistance to fruit tree canker in apple, were investigated using RNA-Seq analysis (Paper V).

3.6.1 Experimental procedures

Trees of two cultivars, the partially resistant ‘Jonathan’ and the susceptible ‘Prima’, were inoculated with *N. ditissima* as previously described (Paper III) and with water as a control. Tissue samples were taken at three different time points, i.e., 5, 15, and 30 dai. In total, 36 tissue samples from the two different apple cultivars at three different time points, each with two inoculations (*N. ditissima* and water) in three replicates (tree), were used for RNA extraction. Concentration, purity and integrity of RNAs were determined, and libraries were generated with 1 µg of total RNA using Illumina TruSeq Stranded mRNA, multiplexed and sequenced in two lanes on Illumina HiSeq High Output mode.

3.6.2 Data analysis

Quality of the reads was determined with the fastQC software version 0.11.2 (Andrews, 2010). The adapter sequences were removed with Trimmomatic, version 0.32 (Bolger & Giorgi, 2014) together with low quality sequences. In order to remove the fungal reads, the reads were mapped to the reference genome of apple, assembly version 1 (Velasco *et al.*, 2010), as well as to the reference genome of the fungus *Nectria haematococca* (Coleman *et al.*, 2009) using TopHat2, version 2.0.13 (Trapnell *et al.*, 2009). A very limited

number of identified fungal reads were removed. Then the filtered reads were used for further analysis. HTseq-count, version 0.6.1 (Anders *et al.*, 2014), was used for calculation of read counts per gene. Normalization and differential expression analysis were performed according to the DESeq2 model and package (Love *et al.*, 2013; Anders & Huber, 2010). A threshold for identification of significantly differentially expressed genes in each cultivar was based on setting the false discovery rate (FDR) to ≤ 0.05 . More details about data analysis are provided in Paper V.

4 Results and discussion

4.1 Variation among apple cultivars inoculated with spores of *N. ditissima*

In the first study (Paper I), damage caused by canker lesions on cut shoots inoculated with *N. ditissima*, was quantified using three different variables: incubation period, infection percentage and AUDPC. Variation among cultivars for each of these variables was determined with two-way ANOVA (Table 1 in Paper I). Significant differentiation among cultivars was obtained with infection percentage and AUDPC, while incubation period produced less informative data and was not used for further analyses. When different replicates were compared, AUDPC values produced the most consistent data (Figure 1 and Table 2 in Paper I) and were therefore used in subsequent analyses.

Significant differences among the 30 inoculated cultivars were found in both years (2012; $P < 0.001$, and 2013; $P < 0.001$). The most resistant cultivar was 'Prairifire' in both years while 'Jonathan' and 'Rödluvan' showed the highest susceptibility in 2012 and 2013, respectively (Figure 11). Additionally, 'Aroma', 'Golden Delicious', 'Santana' and 'Frösåker' were found to be relatively resistant while 'Elise' and 'Hornö' were relatively susceptible in both years. A significant correlation was found across years, 2012 and 2013 ($r = 0.57$, $P < 0.001$). The results for the more resistant cultivars were less varied between years compared to results for the more susceptible cultivars.

In the second study (Paper II), significant differences among cultivars were observed in both sets of experiments, i.e. cut-off shoots and one-year-old potted trees. There was no significant interaction of cultivar \times year for either cut-off shoots or one-year-old trees. Instead a significant correlation was found between data for the two years for each set of experiments, indicating repeatability over years.




Figure 11. Average value for AUDPC for 30 apple cultivars inoculated with *N. ditissima* in the standard greenhouse in 2012 and 2013.

In the cut-off shoots experiment in the biotron, ‘Cox’s Orange Pippin’ was the most susceptible cultivar in both years. The most resistant cultivars were instead ‘Santana’, ‘Golden Delicious’ and ‘Aroma’, and these also had low values for infection percentage. In the one-year-old trees experiment, ‘Golden Delicious’ was the most resistant cultivar in both years, while also ‘Santana’ and ‘Aroma’ were considered as resistant in 2012 and 2013, respectively. ‘Charlamovsky’ was the most susceptible cultivar and also showed the highest value for infection percentage. Additionally ‘Gala’, ‘Cox’s Orange Pippin’, ‘Elise’ and ‘Prima’ were found to be relatively susceptible in both years.

While the cut-off shoots method can produce rough estimates of resistance levels within a short timeframe, inoculation of one year-old trees produces data with superior resolution power and repeatability over years. This method is, however, more expensive and time-consuming and further development of phenotyping procedures is needed. Assessment of infection percentage under ‘natural’ conditions (dispersal of spores onto unwounded potted trees) can be a valuable complement to the wound inoculations, since different components of resistance may be targeted.

The resistance mechanism of apple against fruit tree canker is still not adequately known. Necrotrophic fungi like *N. ditissima*, are usually classified either as host-specific pathogens or as broad host-range pathogens (Mengiste, 2012). The ability of broad-host-range necrotrophs, as *N. ditissima*, to cause disease is usually attributed to the production of a diversity of phytotoxic metabolites, cell wall-degrading enzymes, and cell death elicitors that kill host cells and cause necrosis, whereas host-specific necrotrophs generally produce toxins that have activity on only a limited number of plant species (Laluk & Mengiste, 2010; Friesen *et al.*, 2008). Resistance to necrotrophic fungi is quantitative, and usually requires many genes for full resistance (Laluk & Mengiste, 2010). By contrast, the causal agent of apple scab, *Venturia inaequalis*, is a hemibiotrophic fungus (Jha *et al.*, 2009), and qualitative, race-specific resistance is involved in the defence response of apple to this pathogen including detoxification of fungal toxins, transcriptional regulation and modulation of the defence response (Bastiaanse *et al.*, 2014; Poland *et al.*, 2009).

Previous studies have shown that there is large variation among apple cultivars in level of resistance to *N. ditissima* (Braun, 1997; Pedersen *et al.*, 1994), and different apple cultivars can probably utilize various defence responses to suppress spread and growth of this pathogen. In the current study ‘Prairifire’, which is an ornamental crab apple with a pedigree including *M. atrosanguinea*, *M. zumi calocarpa*, *M. niedzwetskyana* and *M. moerlandsii* ‘Liset’, showed the highest level of resistance to apple canker, which is likely

to be connected with its unique genetic background. Valuable resistance genes have thus been found also in many different *Malus* species. Resistance to the bacterial disease fire blight caused by *Erwinia amylovora*, is controlled by several QTLs with restricted effects in common apple cultivars (Gardiner *et al.*, 2012; Parravicini *et al.*, 2011; Civetta *et al.*, 2009; Flachowsky *et al.*, 2007; Khan *et al.*, 2006), but some major fire blight resistance genes have recently been found in material originating from wild apple species, e.g., *Malus robusta* (Kellerhals *et al.*, 2014; Peil *et al.*, 2007). Several major genes for scab resistance have also been obtained from wild *Malus* species such as *M. floribunda* (Würdig *et al.*, 2015; Belfanti *et al.*, 2004). In keeping with these results, there may be one or several resistance genes in ‘Prairifire’ that could potentially play an important role in future breeding against fruit tree canker.

4.2 Quantitative real-time PCR for detection and quantification of *N. ditissima* in susceptible and resistant apple cultivars

Validation of one primer pair, Bt–fw135/Bt–rw284, showed that a single amplicon (150 bp), representing a specific, targeted sequence of *N. ditissima* gDNA was obtained from *N. ditissima* isolates (Figure 1 in Paper III). Moreover, this primer pair amplified a fragment only in DNA samples of *N. ditissima* and not in samples of any other fungal pathogen (Table 1 in Paper III). Amplification specificity was confirmed by detection of a single specific peak in the qPCR assay. Amplification efficiency of this primer pair was determined to be 99.6% using the standard curve method. An analysis of limit of detection (LOD) showed that the lowest amount of *N. ditissima* gDNA that can be detected by qPCR assay was 1 pg (Figure 12).

Results of the qPCR assay showed that the fungus was not detected in the sampled tissue until two months after inoculation, and then only when samples were taken at the point of infection. The fungus was not detected in the area surrounding the infection site in any of the six studied cultivars.

Variation in the ability to withstand or suppress the development of fungus was observed by qPCR assay in the six cultivars. Samples of the susceptible cultivar ‘Elise’ contained the highest amount of fungal biomass while samples of the partially resistant ‘Aroma’ had the lowest.

In our study, availability of a species-specific primer pair, Bt–fw135/Bt–rw284, derived from the β -tubulin region, allowed detection and quantification of gDNA of *N. ditissima* in the infected tissues. Similarly, qPCR assays with primers designed from the internal transcribed spacer (ITS) regions proved successful in the study of differences between resistant and susceptible

varieties of soybean in terms of accumulation of *Phytophthora sojae* (Catal *et al.*, 2013).

Specific primers have been developed previously from the ITS regions with the aim to detect *N. ditissima* (Langrell, 2002; Langrell & Barbara, 2001), but these primers were not suitable for use in qPCR due to large size of the amplified fragment (412 bp) and cross reactions with apple DNA.

In order to increase the sensitivity of our primer, other approaches like TaqMan-based qPCR can be investigated. As previously shown, the application of SYBR Green primers and TaqMan probe can increase the sensitivity of primers substantially (Garces *et al.*, 2014). It has also been reported that the simultaneous use of two TaqMan probes labeled with two different reporter dyes can increase the assay sensitivity and reproducibility at the low detection range (Yip *et al.*, 2005).




Figure 12. Detection limit of qPCR. Signals 1-6, the results of qPCR for 10 ng, 1 ng, 0.1 ng, 0.01 ng, 1 pg and 0.1 pg DNA per reaction, respectively.

4.3 Histopathology of infection and colonization of susceptible and resistant apple cultivars by *N. ditissima*

For this study, potted trees and cut shoots of the susceptible ‘Cox’s Orange Pippin’ and the partially resistant ‘Santana’, were inoculated with spores of *N. ditissima*. Woody tissue from the affected area was then examined to identify possible differences that could be related to level of susceptibility. Plastic embedding with long incubation times of the specimens resulted in good quality sections from both infected and non-infected samples.

The structural organization of the bark, wood and pith was clearly seen in the transverse, healthy stem sections stained with toluidine blue (Figure 13) and fungal hyphae were visualized by dark purple colour. In one-year-old trees, a tangled mass of fungal hyphae was seen under the periderm of both cultivars at 110 dai, which may be the first step towards the production of sporodochia (Figure 14A). This finding is in agreement with a previous study on the canker of aspen caused by *N. ditissima* (Zalasky, 1968). A larger number of hyphae was observed in cortex parenchyma cells of the susceptible cultivar compared to the partially resistant cultivar, and these cells were completely collapsed in the susceptible cultivar (Figure 14B). Primary and secondary phloem (Figure 14C), primary and secondary xylem, xylem rays and pith cells were affected in the infected tissues of both cultivars (Figure 4 in Paper III). These findings are in agreement with previous studies on apple canker caused by *N. ditissima* (Crowdy, 1949) and infection of *Fraxinus* by *N. ditissima* (Sakamoto et al., 2004). The hyphae could clearly penetrate the secondary cell walls in the phloem fibers, xylem cells and lignified walls, as also reported in a study using ‘Cox’s Orange Pippin’ and ‘Worcester Pearmain’ (Crowdy, 1949). Gel formation was observed in both of our cultivars with no difference in extent or frequency. In summary, ‘Cox’s Orange Pippin’ (susceptible cultivar) was more heavily colonized with *N. ditissima* compared to ‘Santana’, which is a partially resistant cultivar.




Figure13. Transverse section of healthy stem with toluidine blue staining. Overview of the stem showing cortex, phloem, xylem, xylem ray and pith cells. The diameter of the section is 3.5 mm.

In cut shoots, fungal hyphae were seen in cortex, phloem and xylem cells, and phloem cells had collapsed in ‘Cox’s Orange Pippin’ by 18 dai. Cortex and phloem cells had deformed and gel formation was observed in both cultivars at

29 dai. In the resistant cultivar ‘Santana’, fungal hyphae were detected by 36 dai (Figure 5 in Paper III). The pace of infection development differed between the cultivars; fungal hyphae were observed earlier in the susceptible cultivar compared to in the partially resistant one. However, more research is needed to determine the nature of defence mechanisms used against the pathogen in the resistant cultivar.




Figure 14. Transverse section of one-year-old tree 110 days after inoculation in infected cultivar, ‘Cox’s Orange Pippin’. A: A hyphal aggregation that possibly shows an early stage of sporodochial formation. B: Fungal hyphae in cortex parenchyma cells, which were deformed and collapsed. C: Fungal hyphae in secondary phloem, they penetrate through the thick cell wall of fiber cells in the secondary phloem. Scale bar: 20 μm .

Disease progression was slower in one-year-old trees compared with cut shoots. Hyphae were less frequent on the side opposite to the necrotic tissue, in accordance with previous studies (Sakamoto *et al.*, 2004; Crowdy, 1949) where the infection was reported to be relatively local. Some occasional gel formation

was found in both cultivars but no tyloses were detected. To sum up, no significant structural differences were found between resistant and susceptible cultivars, but the spreading of the infection was slower in the partially resistant cultivar. This finding is consistent with previous studies on grapes (Baccari & Lindow, 2011; Chatelet *et al.*, 2011) where no significant differences were found in overall stem anatomy between resistant and susceptible varieties infected by *Xylella fastidiosa*.

The major difference between the susceptible and the resistant cultivar was the degree of colonization by the pathogen. Possibly, a differential release of inhibitors such as phytoalexins that are natural plant antibiotics could be involved in restricting the spread of the pathogen more efficiently in resistant cultivars. Previous studies have shown that degree of host resistance and the level of phytoalexin accumulation are often strongly correlated (Hall *et al.*, 2011; Shi *et al.*, 1991). Therefore, more research is needed to determine whether phytoalexins influence the defence reaction against *Neonectria*, possibly reducing the rate of colonization and disrupting pathogen metabolism.

4.4 Genetic diversity and genetic relationships of *N. ditissima*

SSR primers produced a higher level of band polymorphism across the studied isolates than AFLP primers. This is likely due to detection of multiple alleles at a given locus by SSR markers, whereas AFLP markers mainly detect single alleles at multiple loci randomly distributed in the genome. An AFLP-based analysis of molecular variation (AMOVA) among 44 isolates, obtained from eight different orchards, revealed relatively low genetic differentiation among orchards (11% compared to 89% within-orchard variation).

A principal coordinate analysis (PCoA) was performed with SSR data for 28 isolates (Figure 1 in Paper IV). The isolates were loosely grouped and there was no association with the geographic origin. Similar results were obtained with a PCoA using AFLP data for the same 28 isolates, and with a UPGMA-based cluster analysis using AFLP data for 44 isolates (Figure 2 in Paper IV). In the latter analysis, isolates from the same tree clustered together while isolates from different trees in the same or different orchards did not show any spatial pattern. Gene flow between populations (orchards) may prevent geographic subdivision (Lenormand, 2002), either due to relatively high levels of historical gene flow or high levels of spontaneous mutations and/or to the movement of plant material from one region to another (Carlile *et al.*, 2001).

To study the mating system of this pathogen, single-ascospore isolates from the same perithecium or from different perithecia on the same or different trees, were analysed with both SSR and AFLP markers. The SSR results

suggested that all single-ascospores from a single tree were identical. Considerably more variation was detected with AFLP markers, even among isolates from the same perithecium (Figure 15). Obviously, the fungus is heterothallic allowing outcrossing, but still selfing cannot be ruled out. Previous studies based on spore morphology of *N. ditissima* have reported heterothallic reproduction (Krüger, 1974) as well as homothallic (El-Gholl *et al.*, 1986). The congeneric species *N. haematococca* has both homothallic and heterothallic populations (Coleman, 2008), and this may be true also for *N. ditissima*.

Management of sexually reproducing pathogens is more difficult due to the constant appearance of new genotypes, which increase the variation of features like higher aggressiveness, fungicide resistance, and better fitness in the population. An outcrossing mating system in *N. ditissima* can thus be expected




Figure 15. Dendrogram derived from AFLP band patterns of single-ascospore isolates arising from different perithecia of *N. ditissima*. R: Red Delicious, G: Gravensteiner, H: Holsteiner Cox, P1-3: different perithecia, A1-3: different single-ascospores.

to speed up the evolution of adaptation to resistant hosts and commonly applied fungicides (McDonald & McDermott, 1993).

When the two marker types were compared with a Mantel test, genetic distances among isolates showed no correspondence, possibly due mainly to insufficient number of bands detected with the SSR primers. AFLP markers therefore seem to have a better potential for evaluating genetic diversity and relationships of this pathogen.

4.5 Identification of differentially expressed genes in partially resistant and susceptible cultivars in response to inoculation with *N. ditissima*

In order to identify genes that are differentially expressed in response to the pathogen attack, one partially resistant ('Jonathan') and one susceptible cultivar ('Prima') were investigated, at three different time points after inoculating one-year-old potted trees with spores of *N. ditissima*. A total of 36 RNA samples were collected for RNA-Seq. Approximately 392 million paired end reads were obtained. A total of 345 million Illumina reads passed quality filtering. The majority of reads were mapped to the apple reference genome sequence (Velasco et al., 2010).

A principal component analysis (PCA) of DESeq2 normalized counts for 46157 genes (out of total of 63541) is shown in Figure 16. The lowest number of differentially expressed genes (DEGs) was obtained at 5 dai whereas the highest number was found at 30 dai (Table 1 in Paper V).

A total of 1055 genes, which were differently regulated in the resistant cultivar compared to in the susceptible cultivar, were selected for further analysis. Blasting these DEGs against the KEGG (Kyoto Encyclopedia of Genes and Genomes) database indicate that they are involved in 72 metabolic pathways (Table 3 in Paper V).

A functional classification was carried out first for each cultivar separately using the DEGs (Figure 3 in Paper V) and then with the GO terms of the 1055 gene list (Figure 17). Based on these results, the molecular function group was divided into 11 subcategories, the cellular components into 6 subcategories and the biological processes into 12 subcategories in both analyses.

Several genes were associated with apoptosis processes such as *RPM1*, and some were involved in defense response, e.g., *MLO12* (mildew resistance locus O 12), *Bet v I* (pathogenesis-related proteins), *NPR3* (involved in response to wounding), *ACT7* (involved in response to wounding) and *BAK1* (Leu-rich receptor serine/threonine protein kinase). All of these genes were up-regulated in the partially resistant cultivar 'Jonathan'. The *Bet v I* gene belongs to family

10 of plant pathogenesis-related proteins (PR-10). Pathogenesis-related (PR) proteins have been reported to be induced in plants in the event of a pathogen attack (Linthorst & Van Loon, 1991; Van Loon, 1985).

Several genes involved in the electron transport pathway were also up-regulated in the partially resistant cultivar, e.g., *CYP71A25* (cytochrome P450), *CYP71A26* (encoding oxygen binding) and *ATBCB* (copper binding protein). Cytochromes P450 (CYPs) belong to a superfamily family of proteins and are involved in catabolism of secondary metabolites e.g., hormone and signaling molecules, antioxidants, allelochemicals and defense compounds (Bak *et al.*, 2011; Morant *et al.*, 2003). In this study, the expression level of *CYP71A25* and *CYP71A26* were up-regulated in response to *N. ditissima* inoculation of the partially resistant cultivar. Trognitz *et al.* (2002) showed that cytochromes P450 play a role in the defense response in plants.

Some genes were involved in the flavonoids biosynthesis pathway e.g., *DMR6* (encoding oxidoreductase activity) and *TT10* (Laccase-like polyphenol oxidases). Some genes involved in metabolic processes e.g., *ELI3-1* (elicitor-activated gene 3-1), *GT/UGT74F2* (UDP-glucosyltransferase 74F2), *ATGSTF13* (glutathione transferase) and *CAD1* (glutathione gamma-glutamylcysteinyltransferase 1) were differentially regulated between resistant and susceptible cultivars (Table 2 in Paper V). The *PAL1* and *PAL2* genes, which were up-regulated in the partially resistant cultivar, participate in the defense of plants against pathogens and catalyzes the first step in the biosynthesis of phenylpropanoids, which is responsible for the production of lignin (Dixon *et al.*, 2002). Since an increased activity of the *PAL* gene has been observed in apple leaves infected by *Venturia inaequalis* (Petkovsek *et al.*, 2011; Schovankova & Opatova, 2011), increased *PAL* activity will most likely contribute to reduce also infection by *N. ditissima*.

Expression of some genes such as *WRKY72*, *WRKY70*, *WRKY40*, *MYB2* and transcription factor genes, were induced in response to *N. ditissima* inoculation. The *WRKY* proteins are a large family of transcription factors (Eulgem *et al.*, 2000). *WRKY* family members appear to be involved in the regulation of different physiological programs in plants such as pathogen defense (Eulgem & Somssich, 2007). Eulgem *et al.* (2000) also reported that infection with viruses, bacteria and fungi increased the level of *WRKY* mRNA.

Cytochromes P450 (CYPs) belong to the largest family of plant proteins and are involved in detoxification and biosynthesis of secondary metabolites (Morant *et al.*, 2003), and can play a role in the defense response of plants (Trognitz *et al.*, 2002). The current study demonstrated that expression levels of *CYP71A25* and *CYP71A26* were up-regulated in response to *N. ditissima* inoculation in the partially resistant cultivar.




Figure 16. Principal component analysis of 36 samples using DESeq2 package, which normalized counts for 46157 genes (out of total of 63541).

The response of apple cultivars inoculated with *N. ditissima* also involves the up-regulation of genes involved in fortification of cell walls, detoxification, phosphorylation and pathogen defense.

The results of this study revealed differentially expressed genes at transcription level between a susceptible and a resistant apple cultivar, infected by *N. ditissima*, by utilizing RNA-Seq technology. In addition, it provided a wealth of information for further investigation in order to understand the pathogenicity mechanisms of different apple cultivars against fruit tree canker.




Figure 17. Gene ontology classification with cut off level 2 of differentially expressed genes.

5 Conclusions and future perspectives

The studies conducted within the framework of this thesis can be summarized in the following conclusions:

- Variation in levels of resistance to canker was observed among the investigated apple cultivars. This variation was estimated by different parameters: area under the disease-progression curve (AUDPC), incubation period and infection percentage. Two types of plant models, cut-off shoot and one-year-old tree were used to verify the results. Cultivar differences in resistance found in this study can be exploited in breeding programs.
- A qPCR assay was successfully developed, making it possible to detect and quantify *Neonectria ditissima* in apple cultivars using a species-specific primer pair. However, this assay is not sufficiently sensitive to detect the fungus in asymptomatic tissue. Since there was a relationship between the *N. ditissima* biomass and lesion size in different cultivars, this qPCR assay may be useful for investigating differences between resistant and susceptible cultivars.
- A light microscopy analysis of two infected apple cultivars showed that the fungal hyphae can penetrate into all cell types i.e., cortex, phloem, xylem and pith cells but the infection proceeds faster in susceptible cultivars resulting in a higher amount of fungal hyphae within a given time period. This indicates that level of resistance is not related to the anatomy of the apple woody tissue; instead other mechanisms, e.g., chemical defence, are more likely to be involved in canker resistance.

- The mating system of *N. ditissima* was investigated by application of AFLP analysis of single-ascospore isolates derived from the same perithecium. The results indicate that this fungus is heterothallic, i.e. outcrossing. However, a few isolates were identical, suggesting that homothallic reproduction may also occur.
- Neither SSR nor AFLP markers could detect a spatial pattern of genetic variation among single-ascospore isolates of *N. ditissima* collected in eight orchards, and differentiation between orchards was also quite restricted. Possibly high levels of gene flow due to long-distance transfer of plant material is one reason for the low differentiation and lack of an isolation-by-distance pattern, but high mutation rates can also play a role.
- A gene expression analysis with RNA-Seq data suggests that many genes involved in detoxification, lignification, phosphorylation and pathogen defense are differentially expressed in resistant and susceptible apple cultivars. Some genes with defense-related function, e.g. *Bet v I*, *MLO12*, *ACT7*, *NPR3* and *BAK1*, were induced in the partially resistant apple cultivar during infection.
- The observed changes in gene expression between the susceptible and resistant cultivars highlight possible candidate genes that may play a role in the partial resistance mechanisms of apple and increase our understanding about the molecular mechanisms that are involved against fruit tree canker.

References

- Agrios, G.N. (1997). *Plant pathology*. 5. ed. London: Academic press.
- Ainsworth, G.C. & Bisby, G.R. (2011). *Dictionary of the Fungi*. 10th. ed. Wallingford, Oxon, UK: CAB International.
- Anders, S. & Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biology*, 11(10), R106.
- Anders, S., Pyl, P.T. & Huber, W. (2014). HTSeq–A Python framework to work with high-throughput sequencing data. *Bioinformatics*, doi:10.1093/bioinformatics/btu638.
- Andrews, S. (2010). FastQC: A quality control tool for high throughput sequence data. <http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/>.
- Baccari, C. & Lindow, S.E. (2011). Assessment of the process of movement of *Xylella fastidiosa* within susceptible and resistant grape cultivars. *Phytopathology*, 101(1), pp. 77-84.
- Bak, S., Beisson, F., Bishop, G., Hamberger, B., Höfer, R., Paquette, S. & Werck-Reichhart, D. (2011). Cytochromes P450. *The Arabidopsis Book/American Society of Plant Biologists*, 9, pp. 1-56.
- Baker, K. & Cook, R.J. (1974). *Biological control of plant pathogens*. p. 433.
- Bakhru, H. (1995). *Foods that heal, The natural way to good health*. Orient Paperbacks. ND. India.
- Balconi, C., Stevanato, P., Motto, M. & Biancardi, E. (2012). Breeding for biotic stress resistance/tolerance in plants. *Crop Production for Agricultural Improvement: Springer*, pp. 57-114.
- Bastiaanse, H., Muhovski, Y., Parisi, O., Paris, R., Mingeot, D. & Lateur, M. (2014). Gene expression profiling by cDNA-AFLP reveals potential candidate genes for partial resistance of 'President Roulin' against *Venturia inaequalis*. *BMC genomics*, 15(1), p. 1043.
- Belfanti, E., Silfverberg-Dilworth, E., Tartarini, S., Patocchi, A., Barbieri, M., Zhu, J., Vinatzer, B.A., Gianfranceschi, L., Gessler, C. & Sansavini, S.

- (2004). The *HcrVf2* gene from a wild apple confers scab resistance to a transgenic cultivated variety. *Proceedings of the National Academy of Sciences of the United States of America*, 101(3), pp. 886-890.
- Beltra, R., Ballesteros & Liahos, R. (1969). Studies on the production of growth substances by *Nectria galligena*. *Microbiologia Espanola*, 22(1), pp. 41-54.
- Bénaouf, G. & Parisi, L. (2000). Genetics of host-pathogen relationships between *Venturia inaequalis* races 6 and 7 and *Malus* species. *Phytopathology*, 90(3), pp. 236-242.
- Beresford, R.M. & Kim, K.S. (2011). Identification of regional climatic conditions favorable for development of European canker of apple. *Phytopathology*, 101(1), pp. 135-146.
- Bolger, A. & Giorgi, F. (2014). Trimmomatic: A Flexible Read Trimming Tool for Illumina NGS Data. <http://www.usadellab.org/cms/index.php>.
- Botton, A., Lezzer, P., Dorigoni, A., Barcaccia, G., Ruperti, B. & Ramina, A. (2008). Genetic and environmental factors affecting allergen-related gene expression in apple fruit (*Malus domestica* L. Borkh). *Journal of Agricultural and Food Chemistry*, 56(15), pp. 6707-6716.
- Brandt, R.W. (1964). *Nectria canker of hardwoods*. USDA Forest Service.
- Braun, G. & Craig, B. (2008). Organic apple production guide for atlantic Canada. *Agriculture and Agri-Food Canada*.
- Braun, P. (1997). Distribution and severity of anthracnose canker and European canker of apple in Kings County, Nova Scotia. *Canadian Journal of Plant Pathology*, 19(1), pp. 78-82.
- Carlile, M.J., Watkinson, S.C. & Gooday, G.W. (2001). *The fungi*. Gulf Professional Publishing.
- Castlebury, L.A., Rossman, A.Y. & Hyten, A.S. (2006). Phylogenetic relationships of *Neonectria/Cylindrocarpon* on *Fagus* in North America. *Canadian Journal of Botany*, 84(9), pp. 1417-1433.
- Catal, M., Erler, F., Fulbright, D.W. & Adams, G.C. (2013). Real-time quantitative PCR assays for evaluation of soybean varieties for resistance to the stem and root rot pathogen *Phytophthora sojae*. *European Journal of Plant Pathology*, 137(4), pp. 859-869.
- Cayley, D.M. (1921). Some Observations on the life-history of *Nectria galligena* Bres. *Annals of Botany*, 35(137), pp. 79-92
- Chatelet, D.S., Wistrom, C.M., Purcell, A.H., Rost, T.L. & Matthews, M.A. (2011). Xylem structure of four grape varieties and 12 alternative hosts to the xylem-limited bacterium *Xylella fastidiosa*. *Annals of Botany*, 108(1), pp. 73-85.

- Chaverri, P., Salgado, C., Hirooka, Y., Rossman, A.Y. & Samuels, G.J. (2011). Delimitation of *Neonectria* and *Cylindrocarpon* (Nectriaceae, Hypocreales, Ascomycota) and related genera with *Cylindrocarpon*-like anamorphs. *Studies in Mycology* (68), pp. 57-78.
- Civetta, A., Durel, C.-E., Denancé, C. & Brisset, M.-N. (2009). Two distinct major QTL for resistance to fire blight co-localize on linkage group 12 in apple genotypes 'Evereste' and *Malus floribunda* clone 821. *Genome*, 52(2), pp. 139-147.
- Clatterbuck, W.K. (2006). *Tree Wounds: Response of trees and what you can do*. Agricultural Extension Service: University of Tennessee, UT Extension.
- Coleman, J.J. (2008). *ABC transporters as virulence factors in Nectria haematococca MPVI and genomic analysis of the fungus suggest involvement of horizontal gene transfer in its evolution*. ProQuest.
- Coleman, J.J., Rounsley, S.D., Rodriguez-Carres, M., Kuo, A., Wasmann, C.C., Grimwood, J., Schmutz, J., Taga, M., White, G.J. & Zhou, S. (2009). The genome of *Nectria haematococca*: contribution of supernumerary chromosomes to gene expansion. *PLoS genetics*, 5(8), p. e1000618.
- Cooke, L.R. (1999). The influence of fungicide sprays on infection of apple cv. Bramley's seedling by *Nectria galligena*. *European Journal of Plant Pathology*, 105(8), pp. 783-790.
- Cooke, L.R., Watters, B.S. & Brown, A.E. (1993). The effect of fungicide sprays on the incidence of apple canker (*Nectria galligena*) in bramley seedling. *Plant Pathology*, 42(3), pp. 432-442.
- Crowdy, S.H. (1949). Observations on apple canker .3. The anatomy of the stem canker. *Annals of Applied Biology*, 36(4), pp. 483-495.
- Crowdy, S.H. (1952). Observations on apple canker .4. The infection of leaf scars. *Annals of Applied Biology*, 39(4), pp. 569-580.
- Dewey, F.M. & Swinburne, T.R. (1995). A monoclonal antibody immunoassay for the detection of *Nectria galligena* in apple fruit and woody tissues. *European and Mediterranean Plant Protection Organization*, 25(1-2), pp. 65-73.
- Dixon, R.A., Achnine, L., Kota, P., Liu, C.J., Reddy, M. & Wang, L. (2002). The phenylpropanoid pathway and plant defence-a genomics perspective. *Molecular Plant Pathology*, 3(5), pp. 371-390.
- Dubin, H. & English, H. (1970). Factors affecting apple leaf scar infection by *Nectria galligena* conidia. *Phytopathology*, 64, pp. 1201-1203.
- El-Gholl, N., Barnard, E. & Schroeder, R. (1986). Homothallism in *Nectria galligena*. *Canadian Journal of Botany*, 64(4), pp. 902-903.

- English, H., Dubin, H.J. & Schick, F.J. (1979). Chemical control of European canker of apple. *Plant Disease Reporter*, 63(12), pp. 998-1002.
- Eulgem, T., Rushton, P.J., Robatzek, S. & Somssich, I.E. (2000). The WRKY superfamily of plant transcription factors. *Trends in Plant Science*, 5(5), pp. 199-206.
- Eulgem, T. & Somssich, I.E. (2007). Networks of WRKY transcription factors in defense signaling. *Current opinion in plant biology*, 10(4), pp. 366-371.
- Excoffier, L., Laval, G. & Schneider, S. (2005). Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary bioinformatics online*, 1, p. 47.
- FAO (2013). Food and agriculture organization of the United Nations Statistics Division.
- Farr, D.F., Bills, G.F., Chamuris, G.P. & Rossman, A.Y. (1989). *Fungi on plants and plant products in the United States*. APS press.
- Flachowsky, H., Richter, K., Garcia-Libreros, T., Celton, J.-M., Gardiner, S., Horner, M., Peil, A., Hanke, M.-V. & Bus, V. (2007). Confirmation of the Fire Blight QTL of *Malus × robusta* 5 on Linkage Group 3. *Proceedings of XI International Workshop on Fire Blight 793*, pp. 297-303.
- Flack, N. & Swinburne, T. (1977). Host range of *Nectria galligena* Bres. and the pathogenicity of some Northern Ireland isolates. *Transactions of the British Mycological Society*, 68(2), pp. 185-192.
- Friesen, T.L., Faris, J.D., Solomon, P.S. & Oliver, R.P. (2008). Host-specific toxins: effectors of necrotrophic pathogenicity. *Cellular microbiology*, 10(7), pp. 1421-1428.
- Garces, F.F., Gutierrez, A. & Hoy, J.W. (2014). Detection and quantification of *Xanthomonas albilineans* by qPCR and potential characterization of sugarcane resistance to leaf scald. *Plant Disease*, 98(1), pp. 121-126.
- Gardiner, S.E., Norelli, J.L., de Silva, N., Fazio, G., Peil, A., Malnoy, M., Horner, M., Bowatte, D., Carlisle, C. & Wiedow, C. (2012). Putative resistance gene markers associated with quantitative trait loci for fire blight resistance in *Malus* 'Robusta 5' accessions. *BMC genetics*, 13(1), p. 25.
- Garkava-Gustavsson, L., Zborowska, A., Sehic, J., Rur, M., Nybom, H., Englund, J.E., Lateur, M., Van de Weg, E. & Holefors, A. (2013). Screening of apple cultivars for resistance to european canker, *Neonectria ditissima*. *Acta Horticulturae*, 976, pp. 529-536.

- Gelvonauskiene, D., Sasnauskas, A. & Gelvonauskis, B. (2007). The breeding of apple tree resistant to European canker (*Nectria galligena* Bres.). *Sodininkyste ir Darzininkyste*, 26(3), pp. 174-178.
- Ghasemkhani, M. (2012). *Genetic basis for resistance against fruit tree canker in apple*. Introductory paper at the Faculty of Landscape Planning, Horticulture and Agricultural Science: Swedish University of Agricultural Sciences.
- Ghasemkhani, M., Liljeroth, E., Sehic, J., Zborowska, A. & Nybom, H. (2015). Cut-off shoots method for estimation of partial resistance in apple cultivars to fruit tree canker caused by *Neonectria ditissima*. *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science*, 65(5), pp. 412-421.
- Goethe, R. (1880). Weitere Mitteilungen über den Krebs der Apfelbäume. *Landwirtschaftliche Jahresbericht*, 2, pp. 837-852.
- Gomori, G. (1955). Preparation of buffers for use in enzyme studies. *Methods in Enzymology*, 1, pp. 138-146.
- Grove, G.G. (1990). Nectria canker, in compendium of apple and pear diseases. (eds, Jones, A. L. and Aldwinckle, H. S.) *American Phytopathological Society Press, St-Paul*. Minnesota, USA, pp. 35-36.
- Hall, C., Heath, R. & Guest, D.I. (2011). Rapid and intense accumulation of terpenoid phytoalexins in infected xylem tissues of cotton (*Gossypium hirsutum*) resistant to *Fusarium oxysporum* f.sp *vasinfectum*. *Physiological and molecular plant pathology*, 76(3-4), pp. 182-188.
- Hancock, J.F. (2008). *Temperate fruit crop breeding: germplasm to genomics*. Springer Science & Business Media.
- Hanlin, R.T. (1971). Morphology of *Nectria haematococca*. *American Journal of Botany*, 58(1), pp. 105-116.
- Harris, S.A., Robinson, J.P. & Juniper, B.E. (2002). Genetic clues to the origin of the apple. *TRENDS in Genetics*, 18(8), pp. 426-430.
- Hartig, R. (1880). De Krebspilz der Laubholz Bäume. *Untersuchungen aus dem Forstbotanisches. Institut zu München*, I, pp. 109-128.
- Jha, G., Thakur, K. & Thakur, P. (2009). The *Venturia* apple pathosystem: pathogenicity mechanisms and plant defense responses. *Journal of Biomedicine and Biotechnology*. doi:10.1155/2009/680160.
- Johnson, R. (1992). Past, present and future opportunities in breeding for disease resistance, with examples from wheat. *Euphytica*, 63(1-2), pp. 3-22.
- Juniper, B., Watkins, R. & Harris, S. The origin of the apple. (1996). *Proceedings of Eucarpia Symposium on Fruit Breeding and Genetics 484*, pp. 27-34.

- Kellerhals, M., Schuetz, S., Baumgartner, I.O., Schaad, J., Kost, T., Brogini, G. & Patocchi, A. (2014). Breeding fire blight resistant apple varieties. *Agrarforschung Schweiz*, 5(10), pp. 414-421.
- Kennel, W. (1963). Zur Pathogenese des Obstbaumkrebses (*Nectria galligena* Bres.) am Apfel. *Gartenbauwissenschaft*, pp. 29-64.
- Khachatourians, G.G. (2002). *Transgenic plants and crops*. New York: Marcel Dekker.
- Khan, M.A., Duffy, B., Gessler, C. & Patocchi, A. (2006). QTL mapping of fire blight resistance in apple. *Molecular Breeding*, 17(4), pp. 299-306.
- Kozlovskaya, A.Z., Kurdyuk, T.P. & Marudo, G.M. (1999). Selection for resistance to fungal diseases in apple. *Acta Horticulturae*, 484, pp. 513-517.
- Krüger, J. (1974). Zur Genetik von *Nectria galligena* Bres. *Journal of Phytopathology*, 79(4), pp. 320-342.
- Laluk, K. & Mengiste, T. (2010). Necrotroph attacks on plants: wanton destruction or covert extortion? *The Arabidopsis Book/American Society of Plant Biologists*, 8.
- Langrell, S.R.H. (2000). *Molecular phylogeny, detection and epidemiology of Nectria galligena*. Bres. the incitant of *Nectria canker on apple*. Diss. PhD thesis: University of London.
- Langrell, S.R.H. (2002). Molecular detection of *Neonectria galligena* (syn. *Nectria galligena*). *Mycological Research*, 106, pp. 280-292.
- Langrell, S.R.H. & Barbara, D.J. (2001). Magnetic capture hybridisation for improved PCR detection of *Nectria galligena* from lignified apple extracts. *Plant Molecular Biology Reporter*, 19(1), pp. 5-11.
- Lateur, M. & Populer, C. (1994). Screening fruit tree genetic resources in Belgium for disease resistance and other desirable characters. *Euphytica*, 77(1-2), pp. 147-153.
- Latorre, B.A., Rioja, M.E., Lillo, C. & Munoz, M. (2002). The effect of temperature and wetness duration on infection and a warning system for European canker (*Nectria galligena*) of apple in Chile. *Crop Protection*, 21(4), pp. 285-291.
- Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, 17(4), pp. 183-189.
- Linthorst, H.J. & Van Loon, L. (1991). Pathogenesis-related proteins of plants. *Critical reviews in plant sciences*, 10(2), pp. 123-150.
- Lortie, M. (1964). Production of perithecia of *Nectria galligena* Bres. in pure culture. *Canadian Journal of Botany*, 42(1), pp. 123-124.

- Lortie, M. & Kuntz, J.E. (1963). Ascospore discharge and conidium release by *Nectria galligena* Bres. under field and laboratory conditions. *Canadian Journal of Botany*, 41(8), pp. 1203-1210.
- Love, M., Anders, S. & Huber, W. (2013). Differential analysis of count data—the DESeq2 package. pp. 1-48.
- Luby, J.J., Ferree, D. & Warrington, I. (2003). Taxonomic classification and brief history. *Apples: botany, production and uses*, pp. 1-14.
- Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer research*, 27, pp. 209-220.
- Marra, R.E. & Corwin, J.A. (2009). Isolation and characterization of codominant markers for the perennial canker fungal pathogen *Neonectria ditissima*. *Molecular Ecology Resources*, 9(3), pp. 906-909.
- McCracken, A.R., Berrie, A., Barbara, D.J., Locke, T., Cooke, L.R., Phelps, K., Swinburne, T.R., Brown, A.E., Ellerker, B. & Langrell, S.R.H. (2003). Relative significance of nursery infections and orchard inoculum in the development and spread of apple canker (*Nectria galligena*) in young orchards. *Plant Pathology*, 52(5), pp. 553-566.
- McDonald, B.A. & McDermott, J.M. (1993). Population Genetics of Plant Pathogenic Fungi. *Bioscience*, 43(5), pp. 311-319.
- Mengiste, T. (2012). Plant immunity to necrotrophs. *Annual Review of Phytopathology*, 50, pp. 267-294.
- Mols, P.J.M. & Boers, J.M. (2001). Comparison of a Canadian and a Dutch strain of the parasitoid *Aphelinus mali* (Hald) (Hym., Aphelinidae) for control of woolly apple aphid *Eriosoma lanigerum* (Hausmann) (Hom., Aphididae) in the Netherlands: a simulation approach. *Journal of Applied Entomology*, 125(5), pp. 255-262.
- Morant, M., Bak, S., Møller, B.L. & Werck-Reichhart, D. (2003). Plant cytochromes P450: tools for pharmacology, plant protection and phytoremediation. *Current opinion in biotechnology*, 14(2), pp. 151-162.
- Munson, R. (1939). Observations on apple canker *Annals of Applied Biology*, 26(3), pp. 440-457.
- O'Rourke, D. (2003). *World production, trade, consumption and economic outlook for apples*. Apples: botany, production and uses: CABI Publishing, 02138, USA.
- Parisi, L., Lespinasse, Y., Guillaumes, J. & Krüger, J. (1993). A new race of *Venturia inaequalis* virulent to apples with resistance due to the Vf gene. *Phytopathology*, 83(5), pp. 533-537.

- Parravicini, G., Gessler, C., Denance, C., Lasserre-Zuber, P., Vergne, E., Brisset, M.N., Patocchi, A., Durel, C.E. & Brogini, G.A. (2011). Identification of serine/threonine kinase and nucleotide-binding site-leucine-rich repeat (NBS-LRR) genes in the fire blight resistance quantitative trait locus of apple cultivar 'Evereste'. *Molecular Plant Pathology*, 12(5), pp. 493-505.
- Pedersen, H., Christensen, J.V. & Hansen, P. (1994). Susceptibility of 15 apple cultivars to apple scab, powdery mildew, canker and mites. *Fruit Varieties Journal*, 48(2), pp. 97-100.
- Peil, A., Garcia-Libreros, T., Richter, K., Trognitz, F., Trognitz, B., Hanke, M.V. & Flachowsky, H. (2007). Strong evidence for a fire blight resistance gene of *Malus robusta* located on linkage group 3. *Plant Breeding*, 126(5), pp. 470-475.
- Petkovsek, M.M., Slatnar, A., Stampar, F. & Veberic, R. (2011). Phenolic compounds in apple leaves after infection with apple scab. *Biologia Plantarum*, 55(4), pp. 725-730.
- Pijut, P.M. (2006). *Planting and care of fine hardwood seedlings: Diseases in hardwood tree plantings*. USDA Forest Service, Northern Research Station: FNR-221. Hardwood tree improvement and regeneration center.
- Plante, F., Hamelin, R.C. & Bernier, L. (2002). A comparative study of genetic diversity of populations of *Nectria galligena* and *N. coccinea* var. *faginata* in North America. *Mycological Research*, 106, pp. 183-193.
- Poland, J.A., Balint-Kurti, P.J., Wisser, R.J., Pratt, R.C. & Nelson, R.J. (2009). Shades of gray: the world of quantitative disease resistance. *Trends in Plant Science*, 14(1), pp. 21-29.
- R Development Core Team (2013). *R: A language and environment for statistical computing*, R Foundation for Statistical Computing. Vienna, Austria. <http://www.R-project.org/>.
- Roelfs, A. (1988). Genetic control of phenotypes in wheat stem rust. *Annual Review of Phytopathology*, 26(1), pp. 351-367.
- Rossmann, A.Y. & Palm-Hernandez, M.E. (2008). Systematics of plant pathogenic fungi: Why it matters. *Plant Disease*, 92(10), pp. 1376-1386.
- Rossmann, A.Y., Samuels, G.J., Rogerson, C.T. & Lowen, R. (1999). Genera of *Bionectriaceae*, *Hypocreaceae* and *Nectriaceae* (Hypocreales, Ascomycetes). *Studies in Mycology* (42), pp. 1-248.
- Sakamoto, Y., Yamada, Y., Sano, Y., Tamai, Y. & Funada, R. (2004). Pathological anatomy of *Nectria* canker on *Fraxinus mandshurica* var. *japonica*. *Iawa Journal*, 25(2), pp. 165-174.

- Sasnauskas, A., Gelvonauskiene, D., Gelvonauskis, B., Bendokas, V. & Baniulis, D. (2006). Resistance to fungal diseases of apple cultivars and hybrids in Lithuania. *Agronomy Research*, 4, pp. 349-352.
- Saure, M. (1961). *Untersuchungen über die Voraussetzungen für ein epidemisches Auftreten des Obstbaumkrebses (Nectria galligena Bres.)*. Diss: Obstbauversuchsanstalt.
- Schovankova, J. & Opatova, H. (2011). Changes in phenols composition and activity of phenylalanine-ammonia lyase in apples after fungal infections. *Horticultural Science*, 38(1), pp. 1-10.
- Shaner, G. & Finney, R.E. (1977). Effect of nitrogen-fertilization on expression of slow-mildewing resistance in Knox wheat. *Phytopathology*, 67(8), pp. 1051-1056.
- Shi, J., Mueller, W. & Beckman, C. (1991). Ultrastructural responses of vessel contact cells in cotton plants resistant or susceptible to infection by *Fusarium oxysporum* f. sp. *vasinfectum*. *Physiological and molecular plant pathology*, 38(3), pp. 211-222.
- Swinburne, T. (1971). The infection of apples, cv. Bramley's Seedling, by *Nectria galligena* Bres. *Annals of Applied Biology*, 68(3), pp. 253-262.
- Swinburne, T., Cartwright, J., Flack, N. & Brown, A.E. (1975). The control of apple canker (*Nectria galligena*) in a young orchard with established infections. *Annals of Applied Biology*, 81(1), pp. 61-73.
- Talukder, Z.I., Tharreau, D. & Price, A.H. (2004). Quantitative trait loci analysis suggests that partial resistance to rice blast is mostly determined by race-specific interactions. *New Phytologist*, 162(1), pp. 197-209.
- Trapnell, C., Pachter, L. & Salzberg, S.L. (2009). TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics*, 25(9), pp. 1105-1111.
- Trognitz, F., Manosalva, P., Gysin, R., Niño-Liu, D., Simon, R., del Rosario Herrera, M., Trognitz, B., Ghislain, M. & Nelson, R. (2002). Plant defense genes associated with quantitative resistance to potato late blight in *Solanum phureja* × dihaploid *S. tuberosum* hybrids. *Molecular Plant-Microbe Interactions*, 15(6), pp. 587-597.
- Van de Weg, E. & Jansen, R. (1989). Breeding for resistance to *Nectria galligena*; differences in resistance between seedling populations. Integrated control of pome fruit diseases. *IOBC Bull*, 12(6), pp. 137-145.
- Van de Weg, W.E. (1987). Note on an inoculation method to infect young apple seedlings with *Nectria galligena* Bres. *Euphytica*, 36(3), pp. 853-854.

- Van de Weg, W.E. (1989). Screening for resistance to *Nectria galligena* Bres in cut shoots of apple *Euphytica*, 42(3), pp. 233-240.
- Van Loon, L. (1985). Pathogenesis-related proteins. *Plant Molecular Biology*, 4(2), pp. 111-116.
- Velasco, R., Zharkikh, A., Affourtit, J., Dhingra, A., Cestaro, A., Kalyanaraman, A., Fontana, P., Bhatnagar, S.K., Troggio, M., Pruss, D., Salvi, S., Pindo, M., Baldi, P., Castelletti, S., Cavaiuolo, M., Coppola, G., Costa, F., Cova, V., Dal Ri, A., Goremykin, V., Komjanc, M., Longhi, S., Magnago, P., Malacarne, G., Malnoy, M., Micheletti, D., Moretto, M., Perazzolli, M., Si-Ammour, A., Vezzulli, S., Zini, E., Eldredge, G., Fitzgerald, L.M., Gutin, N., Lanchbury, J., Macalma, T., Mitchell, J.T., Reid, J., Wardell, B., Kodira, C., Chen, Z., Desany, B., Niazi, F., Palmer, M., Koepke, T., Jiwan, D., Schaeffer, S., Krishnan, V., Wu, C., Chu, V.T., King, S.T., Vick, J., Tao, Q., Mraz, A., Stormo, A., Stormo, K., Bogden, R., Ederle, D., Stella, A., Vecchietti, A., Kater, M.M., Masiero, S., Lasserre, P., Lespinasse, Y., Allan, A.C., Bus, V., Chagne, D., Crowhurst, R.N., Gleave, A.P., Lavezzo, E., Fawcett, J.A., Proost, S., Rouze, P., Sterck, L., Toppo, S., Lazzari, B., Hellens, R.P., Durel, C.-E., Gutin, A., Bumgarner, R.E., Gardiner, S.E., Skolnick, M., Egholm, M., Van de Peer, Y., Salamini, F. & Viola, R. (2010). The genome of the domesticated apple (*Malus x domestica* Borkh.). *Nature Genetics*, 42(10), pp. 833-839.
- Ventura, J., Costa, H., Tatagiba, J. & Naqvi, S. (2004). *Diseases of fruits and vegetables: diagnosis and management*. Kluwer, London.
- Weber, R.W.S. (2014). Biology and control of the apple canker fungus *Neonectria ditissima* (syn. *N. galligena*) from a Northwestern European perspective. *Erwerbs-Obstbau*, 56(3), pp. 95-107.
- Weising, K., Nybom, H., Pfenninger, M., Wolff, K. & Kahl, G. (2005). *DNA fingerprinting in plants: principles, methods, and applications*. CRC press.
- Welch, D. (1934). The range and importance of *Nectria* canker on hardwoods in the northeast. *Journal of Forestry*, 32(9), pp. 997-1002.
- Wessel, H. (1979). Sporenfreysetzung bei *Nectria galligena* Bres. in Erwerbs-Apfelanlagen des Niederelbischen Obstbaugesbietes. *Erwerbsobstbau*, 21, pp. 196-201.
- Wiltshire, S.P. (1921). Studies on the apple canker fungus. I. Leaf scar infection. *Annals of Applied Biology*, 8(3-4), pp. 182-192.
- Würdig, J., Flachowsky, H., Saß, A., Peil, A. & Hanke, M.-V. (2015). Improving resistance of different apple cultivars using the *Rvi6* scab

- resistance gene in a cisgenic approach based on the Flp/*FRT* recombinase system. *Molecular Breeding*, 35(3), pp. 1-18.
- Xu, X.M. & Butt, D.J. (1994). Biology and epidemiology of *Nectria galligena* and an infection warning system. *Norwegian Journal of Agricultural Science* 17, pp. 317-324.
- Xu, X.M., Butt, D.J. & Ridout, M.S. (1998). The effects of inoculum dose, duration of wet period, temperature and wound age on infection by *Nectria galligena* of pruning wounds on apple. *European Journal of Plant Pathology*, 104(5), pp. 511-519.
- Xu, X.M. & Robinson, J.D. (2010). Effects of fruit maturity and wetness on the infection of apple fruit by *Neonectria galligena*. *Plant Pathology*, 59(3), pp. 542-547.
- Yang, Y., Shah, J. & Klessig, D.F. (1997). Signal perception and transduction in plant defense responses. *Genes & Development*, 11(13), pp. 1621-1639.
- Yeh, F.C., Yang, R., Boyle, T.B., Ye, Z. & Mao, J.X. (1997). POPGENE, the user-friendly shareware for population genetic analysis. *Molecular biology and biotechnology centre, University of Alberta, Canada*, 10.
- Yip, S.P., To, S.S.T., Leung, P.H.M., Cheung, T.S., Cheng, P.K.C. & Lim, W.W.L. (2005). Use of dual TaqMan probes to increase the sensitivity of 1-step quantitative reverse transcription-PCR: application to the detection of SARS coronavirus. *Clinical Chemistry*, 51(10), pp. 1885-1888.
- Zalasky, H. (1968). Penetration and initial establishment of *Nectria galligena* in aspen and peachleaf willow. *Canadian Journal of Botany*, 46(1), pp. 57-60.
- Zeller, S.M. (1926). European canker of pomaceous fruit trees. *Oregon Agricultural College Experiment Station Experiment Station*, pp. 1-50.

Acknowledgements

I am so grateful to God that gave me the strength to keep going and finish this extremely challenging experience. This has been an exciting journey, several things happened, good, bad and sometimes really bad! Now, it's time to say thank you to all those people who participated in my PhD studies.

I would like to pay special thankfulness, appreciation and warmth to the persons below who made my research successful and assisted me at every point to cherish my goal:

My Supervisor, Professor *Hilde Nybom*, she deserves to be deeply acknowledged for her vital support from amazing writing to invaluable guidance. Her encouragement made it possible to achieve the goal.

My Assistant Supervisors: Dr. *Larisa Gustavsson* for introducing me to apple canker and giving me the opportunity to be a part of this study. Dr. *Salla Marttila* for her fruitful discussions and excellent help with microscopy techniques. Professor *Erland Liljeroth* for his kindness and helping me to solve problems and providing me with insightful suggestions.

Dr. *Erik Alexandersson*, Dr. *Anna Holefors*, Dr. *Kerry Everett*, Dr. *Tetyana Zhebentyayeva*, Dr. *Reiny Scheper* and Dr. *Jonathan Rees-George* for all their theoretical and technical helps.

Dr. *Kristina Rumpunen* for her help in medical issues and gracious hospitality.

Dr. *Björn Canbäck* for his help with bioinformatics analysis.

Dr. *Jan-Eric Englund* for his statistical advice.

Jasna Sehic, Anna Zborowska and Kerstin Brismar for their technical and lab work assistance.

Anna-Sofie Fält, Mari-Louise Skoog and Pia Ohlsson for their sympathetic attitude and being such a good friend and office mate.

Hélen Lindgren and Annelie Ahlman for their energetic smile, unstinted support and consideration.

Professor *Tomas Bryngelsson* and *Åsa Olsson* for great support and welcoming to Alnarp.

Dr. *Mariette Andersson*, Dr. *Helena Persson Hovmalm*, Professor *Anders Carlsson* and Professor *Eva Johansson*, for their help and advice in solving the problems at the uncomfortable moments.

My best friends *Faiza Rasheed, Yanrong Lv, Lijie Zhong, Faraz Muneer* and his wife to celebrate an unforgettable birthday in hospital and all those cheerful moments.

All past and present PhD students for sharing events and experiences.

All the faculty and staff members of Plant Breeding Department at Balsgård and Alnarp, whose services turned my research a success.

And finally:

I would like express my heartfelt thanks to my husband, who always surprises me with his kindness even in the most difficult times, and to my Mom and Dad. Thanks for your love, patience, support morally and emotionally and encouragement.

You are always my biggest motivation to overcome any obstacles that I encountered.